

**City: North Arlington** 

# VACCINATION INTAKE FORM

			Patio	ent In	format	ion					
Patient Name:										Date:	
Patient Address:											
City:		State	:	Zip:			DOB:			Gender: □ Ma	ale 🗆 Female
Phone #:		Allerg	jies:								
Alt Contact:		Relati	ionship:							Phone:	
Weight (select on	ne): □ < 130 lbs □ 1	31-15	2 lbs □ 1	153-20	0 lbs	□ 200	)-259	lbs [	□ > 260	) lbs	
НСРС	Product		DX	Dos	sage	Inje	ction	Site		Lot#	Exp.
90658 IIV	<i>l</i> :	® 	<b>Z23</b>	0.5	i mL	L/R	DELT	OID IM			
	Boostrix ®		<b>Z23</b>	0.5	i mL	L/R	DELT	OID IM			
	Shingrix ®		<b>Z23</b>	0.5	i mL	L / R DELTOID IM					
	Prevnar ®		<b>Z23</b>	0.5	i mL	L / R DELT		OID IM			
	Pneumovax ®		<b>Z23</b>	0.5	i mL	L / R DELTOID IM					
	Havrix 1440 unit ®		<b>Z23</b>	1 mL		L / R DELTOID IM					
			Physi	cian I	nforma	ition					
Ordering Physicia	an: Eric Jackson				Physic	an Pho	ne:				
NPI: 1235131251	1			Physician Fax:							
Address: 166-18	Ridge Road										
City: North Arlington			State: NJ				Zip: 07031				
			Adminis	sterin	g Phar	macis	t				
Name:											
Address: 98 Ridg	ge Road										

State: NJ

Zip: 07031



### **HB Pharmacy Vaccination Consent Form**

I have read or had explained to me the Vaccine Information Statementforthe current influenza vaccine and understand the risk and benefit.

I understand that receipt of the vaccine does not completely protect me against the flu or other illnesses that resemble the flu. I further understand that if I have a condition (or am undergoing treatment which causes) or immune-suppression (the reduction in my body's ability to fight infection and illness), the effectiveness of the vaccine in preventing the flu may be diminished.

I GIVE CONSENT to HB Pharmacy and licensed staff fo administer this vaccine to me. I hereby release HB Pharmacy and associated staff from any liability for giving me the influenza vaccination. I agree to defend and hold HB Pharmacy and associated staff harmless from any claim made by any person.

My signature on this form means that all of the information provided to HB Pharmacy is true to the best of my knowledge. If this consent form is not signed, dated and returned I will not be vaccinated.

Patient Signature:	Date:
Patient Name (Print):	
□ I authorize HB Pharmacy to fax a Notification of Vacci	nation Letter to my Primary Care Provider:
Physician's Name:	Fax Number:
$\square$ I do not authorize HB Pharmacy to fax a Notification o Provider.	of Vaccination letter to niy Primary Care
*If neither box is checked, a Notification of Vaccination Letter will	be faxed to my Primary Care Provider, if identified.
VIS Given:	Version Date:

New Jersey Department of Health Vaccine Preventable Disease Program P.O. Box 369, Trenton, NJ 08625-0369 609-826-4860 (Fax 609-826-4866) www.njiis.nj.gov

### NEW JERSEY IMMUNIZATION INFORMATION SYSTEM (NJIIS) CONSENT TO PARTICIPATE

#### - RETAIN A COPY OF THIS FORM IN THE MEDICAL RECORD -

	Т					
REGISTRANT INFORMATION	PARENT/GUARDIAN INFORMATION (if NJIIS Registrant is a minor)					
Registrant Name (Print)	Name (Print)					
Date of Birth	Address					
Country of Birth	City, State, Zip Code					
Name of Primary Health Care Provider	Relationship to Registrant					
I have received information about the New Jersey Immunization of this program is to help remind me when my/my child's imchild's immunization history.  I understand that the medical information in the NJIIS may licensed child care centers, colleges, public health agencies,	nmunizations are due and to keep y be shared with authorized hea health insurance companies, and	p a central record of my/my  alth care providers, schools,				
Jersey Law at N.J.S.A. 26:4-131 et seq. and rules at N.J.A.C. I understand that I can get a copy of my/my child's record from or the New Jersey Department of Health (NJDOH). The Nulisted above.	n my primary health care provider,					
There is no cost to participate in this program.						
Yes, I would like to participate in this program.						
□No, I do not want to participate in this program.						
Signature of Registrant (or Parent/Guardian, IF Registrant und	der 18 Years of Age) Date					
	•					
Name of NJIIS Enrollment Site	Registry ID Number	Medical Record Number				

# Screening Checklist for Contraindications to Vaccines for Adults

ATIENT NAME	
ATE OF BIRTH/	
monun day year	

For patients: The following questions will help us determine which vaccines you may be given today. If you answer "yes" to any question, it does not necessarily mean you should not be vaccinated. It just means we need to ask you more questions. If a question is not clear, please ask your healthcare provider to explain it.

		yes	no	know
1. Are you sic	k today?			
2. Do you hav	re allergies to medications, food, a vaccine ingredient, or latex?			
3. Have you e	ver had a serious reaction after receiving a vaccine?			
(e.g., diabet	re a long-term health problem with heart, lung, kidney, or metabolic disease res), asthma, a blood clotting disorder, no spleen, complement component deficiency, mplant, or a spinal fluid leak? Are you on long-term aspirin therapy?			
5. Do you hav	re cancer, leukemia, HIV/AIDS, or any other immune system problem?			
6. Do you hav	re a parent, brother, or sister with an immune system problem?			
as predniso	3 months, have you taken medicines that affect your immune system, such ne, other steroids, or anticancer drugs; drugs for the treatment of rheumatoid ohn's disease, or psoriasis; or have you had radiation treatments?			
8. Have you h	ad a seizure or a brain or other nervous system problem?			
_	past year, have you received a transfusion of blood or blood products, en immune (gamma) globulin or an antiviral drug?			
<b>10.</b> Are you pre	egnant or is there a chance you could become pregnant during the next month?			
<b>11.</b> Have you r	eceived any vaccinations in the past 4 weeks?			
	FORM COMPLETED BY	DATE		
	FORM REVIEWED BY	DATE		
	Did you bring your immunization record card with you?  It is important for you to have a personal record of your vaccinations. If you don't ha ask your healthcare provider to give you one. Keep this record in a safe place and bring seek medical care. Make sure your healthcare provider records all your vaccinations.	g it with yοι		





# Screening Checklist for Contraindications DATE OF BIRTH Month / day / year to Vaccines for Children and Teens

For parents/guardians: The following questions will help us determine which vaccines your child may be given today. If you answer "yes" to any question, it does not necessarily mean your child should not be vaccinated. It just means additional questions must be asked. If a question is not clear, please ask your healthcare provider to explain it.

		yes	110	know
1. Is the child	d sick today?			
2. Does the	child have allergies to medications, food, a vaccine component, or latex?			
3. Has the ch	hild had a serious reaction to a vaccine in the past?			
(e.g., diab	child have a long-term health problem with lung, heart, kidney or metabolic disease etes), asthma, a blood disorder, no spleen, complement component deficiency, implant, or a spinal fluid leak? Is he/she on long-term aspirin therapy?			
	to be vaccinated is 2 through 4 years of age, has a healthcare provider told you nild had wheezing or asthma in the past 12 months?			
<b>6.</b> If your chi	ld is a baby, have you ever been told he or she has had intussusception?			
	hild, a sibling, or a parent had a seizure; has the child had brain or other ystem problems?			
8. Does the	child have cancer, leukemia, HIV/AIDS, or any other immune system problem?			
9. Does the	child have a parent, brother, or sister with an immune system problem?			
as prednis	t 3 months, has the child taken medications that affect the immune system such sone, other steroids, or anticancer drugs; drugs for the treatment of rheumatoid Crohn's disease, or psoriasis; or had radiation treatments?			
	t year, has the child received a transfusion of blood or blood products, or been nune (gamma) globulin or an antiviral drug?			
<b>12.</b> Is the child next mont	d/teen pregnant or is there a chance she could become pregnant during the			
13. Has the ch	hild received vaccinations in the past 4 weeks?			
	FORM COMPLETED BY	DATE		
	FORM REVIEWED BY	DATE.		
	Did you bring your immunization record card with you? yes $\Box$ no $\Box$ It is important to have a personal record of your child's vaccinations. If you don't have			
immunization	healthcare provider to give you one with all your child's vaccinations on it. Keep it is it with you every time you seek medical care for your child. Your child will need this care or school, for employment, or for international travel.			



yes

## **Notification of Vaccination Letter Template**

Dear doctor or nurse at	CARE CLINIC	
We recently pr have informati	ovided vaccination services to your patien on about the vaccines we administered so contact us if you have any questions abo	o you can update your patient's medical
	d the patient (or parent/guardian) with a w	
☐ We entered	I information about the vaccine(s) we adrion information system.	(,,
		(MM/DD/YR)
(For a child, parent/guardian name	Parent,	/guardian birthdate(MM/DD/YR)
The vaccine(s) we administered on	is/are checked below	
VACCINES ADMINISTERED		
COVID-19    mRNA (circle one): Moderna Pfizer   viral vector (Janssen [Johnson &	Hib (monovalent)  ☐ ActHIB (PRP-T)  ☐ Hiberix (PRP-T)	<ul><li>☐ Human papillomavirus (9vHPV) (Gardasil 9)</li><li>☐ MMR</li></ul>
Johnson])	☐ PedvaxHIB (PRP-OMP)	☐ <b>Varicella</b> (chickenpox) (Varivax)
Hepatitis B	☐ Influenza	☐ <b>MMRV</b> (ProQuad)
<ul> <li>Engerix-B; Recombivax HB;</li> <li>DOSE (circle one): 0.5 mL 1.0 mL</li> <li>Heplisav-B (age 18 yrs and older)</li> </ul>	BRAND  DOSE (mL)  ROUTE (circle one): IM Nasal	☐ Hepatitis A (Havrix; Vaqta)  ☐ DOSE (circle one): 0.5 mL 1.0 mL  ☐ HepA-HepB (Twinrix)
<ul> <li>□ PreHevbrio (age 18 yrs and older)</li> <li>□ DTaP (age 6 yrs and younger)</li> <li>□ DTaP-HepB-IPV (Pediarix)</li> </ul>	☐ IPV (Polio)  Pneumococcal conjugate (PCV)	☐ Meningococcal ACWY (MenACWY) (circle one): (Menactra, MenQuadfi, Menveo)
□ DTaP-IPV (Kinrix, Quadracel) □ DTaP-IPV/Hib (Pentacel)	<ul><li>☐ PCV13, Prevnar 13 (through age 18 yrs)</li><li>☐ PCV15, Vaxneuvance</li><li>☐ PCV20, Prevnar 20</li></ul>	Meningococcal B (MenB)  ☐ Bexsero (MenB-4C)
<ul><li>□ DTaP-IPV-Hib-HepB (Vaxelis)</li><li>□ DT (through age 6 yrs)</li></ul>	☐ Pneumococcal polysaccharide (PPSV23) (Pneumovax 23)	<ul><li>☐ Trumenba (MenB-FHbp)</li><li>☐ Zoster (shingles) (RZV) (Shingrix)</li></ul>
<ul><li>☐ <b>Tdap</b> (age 7 yrs and older)</li><li>☐ <b>Td</b> (age 7 yrs and older)</li></ul>	Rotavirus  ☐ RV1 (Rotarix) ☐ RV5 (RotaTeq)	□ Other
NAME OF CLINIC PROVIDING SERVICES	CLINIC CONTACT PERSON	1
ADDRESS	EMAIL ADDRESS	
CITY/STATE/ZIP	PHONE	



# Vaccine Administration Record for Adults

Patient name:	
Birthdate:	Chart number:
Clinic name and address	

Before administering any vaccines, give the patient copies of all pertinent Vaccine Information Statements (VISs) and make sure he/she understands the risks and benefits of the vaccine(s). Always provide or update the patient's personal record card.

Vaccine	Type of Vaccine <sup>1</sup>	Date given	Funding	Noute	Vaccine		Vaccine In Stateme	Vaccinator <sup>5</sup> (signature or	
	Type or vaccine	(mo/day/yr)	(F,S,P) <sup>2</sup>	& Site <sup>3</sup>	Lot#	Mfr.	Date on VIS <sup>4</sup>	Date given <sup>4</sup>	initials & title)
Tetanus, Diphtheria, Pertussis (e.g., Td, Tdap) Give IM. <sup>3</sup>									
Hepatitis A <sup>6</sup> (e.g., HepA, HepA-HepB) Give IM. <sup>3</sup>									
Hepatitis B <sup>6</sup> (e.g., HepB, HepA-HepB) Give IM. <sup>3</sup>									
Human papillomavirus (HPV2, HPV4) Give IM. <sup>3</sup>									
Measles, Mumps, Rubella (MMR) Give SC. <sup>3</sup>									
Varicella (VAR) Give SC. <sup>3</sup>									
Pneumococcal (e.g., PCV13, conjugate; PPSV23, polysaccharide) Give PCV13 IM. <sup>3</sup>									
Give PPSV23 IM or SC. <sup>3</sup>									
Meningococcal (e.g., MenACWY, conjugate; MPSV4, polysaccharide) Give MenACWY IM. <sup>3</sup>									
Give MenACWY IM. Give MPSV4 SC.3									

See page 2 to record influenza, Hib, zoster, and other vaccines (e.g., travel vaccines).

#### **How to Complete This Record**

- 1. Record the generic abbreviation (e.g., Tdap) or the trade name for each vaccine (see table at right).
- 2. Record the funding source of the vaccine given as either F (federal), S (state), or P (private).
- 3. Record the route by which the vaccine was given as either intramuscular (IM), subcutaneous (SC), intradermal (ID), intranasal (IN), or oral (PO) and also the site where it was administered as either RA (right arm), LA (left arm), RT (right thigh), or LT (left thigh).
- Record the publication date of each VIS as well as the date the VIS is given to the patient.
- 5. To meet the space constraints of this form and federal requirements for documentation, a healthcare setting may want to keep a reference list of vaccinators that includes their initials and titles.
- 6. For combination vaccines, fill in a row for each antigen in the combination.

Abbreviation	Trade Name and Manufacturer						
Tdap	Adacel (sanofi pasteur); Boostrix (GlaxoSmithKline [GSK])						
Td	Decavac (sanofi pasteur); generic Td (MA Biological Labs)						
НерА	Havrix (GSK); Vaqta (Merck)						
НерВ	Engerix-B (GSK); Recombivax HB (Merck)						
НерА-НерВ	Twinrix (GSK)						
HPV2	Cervarix (GSK)						
HPV4	Gardasil (Merck)						
MMR	MMRII (Merck)						
VAR	Varivax (Merck)						
PCV13, PPSV23	Prevnar 13 (Pfizer); Pneumovax 23 (Merck)						
MenACWY	Menactra (sanofi pasteur); Menveo (Novartis)						
MPSV4	Menomune (sanofi pasteur)						

# Vaccine Administration Record for Adults

Patient name:		
Birthdate:	Chart number:	
Clinic name and address		

Before administering any vaccines, give the patient copies of all pertinent Vaccine Information Statements (VISs) and make sure he/she understands the risks and benefits of the vaccine(s). Always provide or update the patient's personal record card.

Vaccine	Type of Vaccine <sup>1</sup>	Date given	Funding Source	Route	Vaccine		Vaccine In Stateme	Vaccinator <sup>5</sup> (signature or	
	.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	(mo/day/yr)	(F,S,P) <sup>2</sup>	& Site <sup>3</sup>	Lot#	Mfr.	Date on VIS <sup>4</sup>	Date given <sup>4</sup>	initials & title)
Influenza									
(e.g., IIV3, trivalent inactivated;									
IIV4, quadrivalent inactivated;									
RIV. recombinant inac-									
tivated; LAIV4, quadrivalent live									
LAIV4, quadrivalent live attenuated) Give IIV and RIV IM. <sup>3</sup>									
Give LAIV IN. <sup>3</sup>									
Hib Give IM. <sup>3</sup>									
Zoster (Zos) Give SC.3									
Other									

See page 1 to record Tdap/Td, hepatitis A, hepatitis B, HPV, MMR, varicella, pneumococcal, and meningococcal vaccines.

#### **How to Complete This Record**

- 1. Record the generic abbreviation (e.g., Tdap) or the trade name for each vaccine (see table at right).
- 2. Record the funding source of the vaccine given as either F (federal), S (state), or P (private).
- 3. Record the route by which the vaccine was given as either intramuscular (IM), subcutaneous (SC), intradermal (ID), intranasal (IN), or oral (PO) and also the site where it was administered as either RA (right arm), LA (left arm), RT (right thigh), or LT (left thigh).
- 4. Record the publication date of each VIS as well as the date the VIS is given to the patient.
- 5. To meet the space constraints of this form and federal requirements for documentation, a healthcare setting may want to keep a reference list of vaccinators that includes their initials and titles.

Abbreviation	Trade Name and Manufacturer
LAIV (Live attenuated influenza vaccine)	FluMist (MedImmune)
IIV (Inactivated influ- enza vaccine), RIV (recombinant influenza vaccine)	Afluria (CSL Biotherapies); Agriflu (Novartis); Fluarix (GSK); Flublok (Protein Sciences Corp.); Flucelvax (Novartis); FluLaval (GSK); Fluvinin (Novartis); Fluzone, Fluzone Intradermal, Fluzone High-Dose (sanofi pasteur)
Hib	ActHIB (sanofi pasteur); Hiberix (GSK); PedvaxHib (Merck)
ZOS (shingles)	Zostavax (Merck)

#### **VACCINE INFORMATION STATEMENT**

# Recombinant Zoster (Shingles) Vaccine: What You Need to Know


Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

#### 1. Why get vaccinated?

Recombinant zoster (shingles) vaccine can prevent shingles.

Shingles (also called herpes zoster, or just zoster) is a painful skin rash, usually with blisters. In addition to the rash, shingles can cause fever, headache, chills, or upset stomach. Rarely, shingles can lead to complications such as pneumonia, hearing problems, blindness, brain inflammation (encephalitis), or death.



The risk of shingles increases with age. The most common complication of shingles is long-term nerve pain called postherpetic neuralgia (PHN). PHN occurs in the areas where the shingles rash was and can last for months or years after the rash goes away. The pain from PHN can be severe and debilitating.

The risk of PHN increases with age. An older adult with shingles is more likely to develop PHN and have longer lasting and more severe pain than a younger person.

People with weakened immune systems also have a higher risk of getting shingles and complications from the disease.

Shingles is caused by varicella-zoster virus, the same virus that causes chickenpox. After you have chickenpox, the virus stays in your body and can cause shingles later in life. Shingles cannot be passed from one person to another, but the virus that causes shingles can spread and cause chickenpox in someone who has never had chickenpox or has never received chickenpox vaccine.

#### 2. Recombinant shingles vaccine

Recombinant shingles vaccine provides strong protection against shingles. By preventing shingles, recombinant shingles vaccine also protects against PHN and other complications.

Recombinant shingles vaccine is recommended for:

- Adults 50 years and older
- Adults 19 years and older who have a weakened immune system because of disease or treatments

Shingles vaccine is given as a two-dose series. For most people, the second dose should be given 2 to 6 months after the first dose. Some people who have or will have a weakened immune system can get the second dose 1 to 2 months after the first dose. Ask your health care provider for guidance.

People who have had shingles in the past and people who have received varicella (chickenpox) vaccine are recommended to get recombinant shingles vaccine. The vaccine is also recommended for people who have already gotten another type of shingles vaccine, the live shingles vaccine. There is no live virus in recombinant shingles vaccine.

Shingles vaccine may be given at the same time as other vaccines.



## 3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an allergic reaction after a previous dose of recombinant shingles vaccine, or has any severe, life-threatening allergies
- Is currently experiencing an episode of shingles
- Is pregnant

In some cases, your health care provider may decide to postpone shingles vaccination until a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting recombinant shingles vaccine.

Your health care provider can give you more information.

#### 4. Risks of a vaccine reaction

- A sore arm with mild or moderate pain is very common after recombinant shingles vaccine.
   Redness and swelling can also happen at the site of the injection.
- Tiredness, muscle pain, headache, shivering, fever, stomach pain, and nausea are common after recombinant shingles vaccine.

These side effects may temporarily prevent a vaccinated person from doing regular activities. Symptoms usually go away on their own in 2 to 3 days. You should still get the second dose of recombinant shingles vaccine even if you had one of these reactions after the first dose.

Guillain-Barré syndrome (GBS), a serious nervous system disorder, has been reported very rarely after recombinant zoster vaccine.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

## 5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at <a href="www.vaers.hhs.gov">www.vaers.hhs.gov</a> or call 1-800-822-7967. VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.

#### 6. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at <a href="https://www.fda.gov/vaccines-blood-biologics/vaccines">www.fda.gov/vaccines-blood-biologics/vaccines</a>.
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call 1-800-232-4636 (1-800-CDC-INFO) or
  - Visit CDC's website at <a href="https://www.cdc.gov/vaccines">www.cdc.gov/vaccines</a>.

